


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 405/00 // A61K 31/557	A1	(11) International Publication Number: WO 99/12898 (43) International Publication Date: 18 March 1999 (18.03.99)
<p>(21) International Application Number: PCT/US98/18594</p> <p>(22) International Filing Date: 4 September 1998 (04.09.98)</p> <p>(30) Priority Data: 60/058,252 9 September 1997 (09.09.97) US</p> <p>(71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).</p> <p>(72) Inventors: DELONG, Mitchell, Anthony; 8084 Tyler's Circle, West Chester, OH 45069 (US). WOS, John, August; 8505 Harperpoint Drive, Cincinnati, OH 45249 (US). AMBURGEY, Jack, S., Jr.; 265 Stockton Drive, Loveland, OH 45140 (US). DE, Biswanath; 11269 Cornell Woods Drive, Cincinnati, OH 45241 (US). DAI, Haiyan, George; 47-2 Revere Road, Drexel Hill, PA 19026 (US). SOPER, David, Lindsey; Apartment #3, 298 Courtland Drive, Monroe, OH 45050 (US).</p> <p>(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).</p>		<p>(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>

(54) Title: AROMATIC C₁₆-C₂₀-SUBSTITUTED TETRAHYDRO PROSTAGLANDINS USEFUL AS FP AGONISTS

(57) Abstract

The invention provides novel PGF analogs. In particular, the present invention relates to compounds having a structure according to formula (I): wherein R₁, R₂, R₃, R₄, X, and Z are as defined. This invention also includes optical isomers, diastereomers and enantiomers of said formula, and pharmaceutically-acceptable salts, biohydrolyzable amides, esters, and imides thereof. The compounds of the present invention are useful for the treatment of a variety of diseases and conditions, such as bone disorders and glaucoma. Accordingly, the invention further provides pharmaceutical compositions comprising these compounds. The invention still further provides methods of treatment for bone disorders and glaucoma using these compounds or the compositions containing them.



(1)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


AROMATIC C₁₆ - C₂₀ - SUBSTITUTED TETRAHYDRO PROSTAGLANDINS
USEFUL AS FP AGONISTS

TECHNICAL FIELD

The subject invention relates to certain novel analogs of the naturally occurring prostaglandins. Specifically, the subject invention relates to novel Prostaglandin F analogs. The subject invention further relates to methods of using said novel Prostaglandin F analogs. Preferred uses include methods of treating bone disorders and glaucoma.

BACKGROUND OF THE INVENTION

Naturally occurring prostaglandins (PGA, PGB, PGE, PGF, and PGI) are C-20 unsaturated fatty acids. PGF_{2α}, the naturally occurring Prostaglandin F in humans, is characterized by hydroxyl groups at the C₉ and C₁₁ positions on the alicyclic ring, a cis-double bond between C₅ and C₆, and a trans-double bond between C₁₃ and C₁₄. Thus PGF_{2α} has the following formula:



Analogues of naturally occurring Prostaglandin F have been disclosed in the art. For example, see U.S. Patent No. 4,024,179 issued to Bindra and Johnson on May 17, 1977; German Patent No. DT-002,460,990 issued to Beck, Lerch, Seeger, and Teufel published on July 1, 1976; U.S. Patent No. 4,128,720 issued to Hayashi, Kori, and Miyake on December 5, 1978; U.S. Patent No. 4,011,262 issued to Hess, Johnson, Bindra, and Schaaf on March 8, 1977; U.S. Patent No. 3,776,938 issued to Bergstrom and Sjovald on December 4, 1973; P.W. Collins and S. W. Djuric, "Synthesis of Therapeutically Useful Prostaglandin and Prostacyclin Analogs", Chem. Rev. Vol. 93 (1993), pp. 1533-1564; G. L. Bundy and F. H. Lincoln, "Synthesis of 17-Phenyl-18,19,20-Trinorprostaglandins: I. The PG₁ Series", Prostaglandins, Vol. 9 No. 1 (1975), pp. 1-4; W. Bartman, G. Beck, U. Lerch, H. Teufel, and B. Scholkens, "Luteolytic

Prostaglandins: Synthesis and Biological Activity", Prostaglandins, Vol. 17 No. 2 (1979), pp. 301-311; C. Liljebris, G. Selen, B. Resul, J. Sternschantz, and U. Hacksell, "Derivatives of 17- Phenyl-18,19,20-trinorprostaglandin F₂ α Isopropyl Ester: Potential Antiglaucoma Agents", Journal of Medicinal Chemistry, Vol. 38 No. 2 (1995), pp. 289-304.


Naturally occurring prostaglandins are known to possess a wide range of pharmacological properties. For example, prostaglandins have been shown to: relax smooth muscle, which results in vasodilatation and bronchodilatation, to inhibit gastric acid secretion, to inhibit platelet aggregation, to reduce intraocular pressure, and to induce labor. Although naturally occurring prostaglandins are characterized by their activity against a particular prostaglandin receptor, they generally are not specific for any one prostaglandin receptor. Therefore, naturally-occurring prostaglandins are known to cause side effects such as inflammation, as well as surface irritation when administered systemically. It is generally believed that the rapid metabolism of the naturally occurring prostaglandins following their release in the body limits some of the effects of the prostaglandin to a local area. This effectively prevents the prostaglandin from stimulating prostaglandin receptors throughout the body and causing the effects seen with the systemic administration of naturally occurring prostaglandins.

Prostaglandins, especially prostaglandins of the E series (PGE), are known to be potent stimulators of bone resorption. PGF_{2 α} has also been shown to be a stimulator of bone resorption but not as potent as PGE₂. Also, it has been demonstrated the PGF_{2 α} has little effect on bone formation. It has been suggested that some of the effects of PGF_{2 α} on bone resorption, formation and cell replication may be mediated by an increase in endogenous PGE₂ production.

In view of both the wide range of pharmacological properties of naturally occurring prostaglandins and of the side effects seen with the systemic administration of these naturally occurring prostaglandins, attempts have been made to prepare analogs to the naturally occurring prostaglandins that are selective for a specific receptor or receptors. A number of such analogs have been disclosed in the art. Though a variety of prostaglandin analogs have been disclosed, there is a continuing need for potent, selective prostaglandin analogs for the treatment of a variety diseases and conditions.

SUMMARY OF THE INVENTION

The invention provides novel PGF analogs. In particular, the present invention relates to compounds having a structure according to the following formula:



wherein R_1 , R_2 , R_3 , R_4 , X , and Z are defined below.

This invention also includes optical isomers, diastereomers and enantiomers of the formula above, and pharmaceutically-acceptable salts, biohydrolyzable amides, esters, and imides thereof.

The compounds of the present invention are useful for the treatment of a variety of diseases and conditions, such as bone disorders and glaucoma. Accordingly, the invention further provides pharmaceutical compositions comprising these compounds. The invention still further provides methods of treatment for bone disorders and glaucoma using these compounds or the compositions containing them.

DETAILED DESCRIPTION OF THE INVENTION

Terms and Definitions

"Acyl" is a group suitable for acylating a nitrogen atom to form an amide or carbamate or an oxygen atom to form an ester group. Preferred acyl groups include benzoyl, acetyl, tert-butyl acetyl, para-phenyl benzoyl, and trifluoroacetyl. More preferred acyl groups include acetyl and benzoyl. The most preferred acyl group is acetyl.

"Alkyl" is a saturated or unsaturated hydrocarbon chain having 1 to 18 carbon atoms, preferably 1 to 12, more preferably 1 to 6, more preferably still 1 to 4 carbon atoms. Alkyl chains may be straight or branched. Preferred branched alkyl have one or two branches, preferably one branch. Preferred alkyl are saturated. Unsaturated alkyl have one or more double bonds and/or one or more triple bonds. Preferred unsaturated alkyl have one or two double bonds or one triple bond, more preferably one double bond. Alkyl chains may be unsubstituted or substituted with from 1 to 4 substituents. Preferred alkyl are unsubstituted. Preferred substituted alkyl are mono-, di-, or trisubstituted. Preferred alkyl substituents include halo, haloalkyl, hydroxy, aryl (e.g., phenyl, tolyl, alkyloxyphenyl, alkyloxycarbonylphenyl, halophenyl), heterocyclyl, and heteroaryl.

"Aromatic ring" is an aromatic hydrocarbon ring system. Aromatic rings are monocyclic or fused bicyclic ring systems. Monocyclic aromatic rings contain from about 5 to about 10 carbon atoms, preferably from 5 to 7 carbon atoms, and most preferably from 5 to 6 carbon atoms in the ring. Bicyclic aromatic rings contain from 8 to 12 carbon atoms, preferably 9 or 10 carbon atoms in the ring. Aromatic rings may be unsubstituted or substituted with from 1 to 4 substituents on the ring. Preferred aromatic ring substituents include: halo, cyano, alkyl, heteroalkyl, haloalkyl, phenyl, phenoxy or any combination thereof. More preferred substituents include halo and haloalkyl. Preferred aromatic rings include naphthyl and phenyl. The most preferred aromatic ring is phenyl.

"Carbocyclic aliphatic ring" is a saturated or unsaturated hydrocarbon ring. Carbocyclic aliphatic rings are not aromatic. Carbocyclic aliphatic rings are monocyclic, or are fused, spiro, or bridged bicyclic ring systems. Monocyclic carbocyclic aliphatic rings contain from about 4 to about 10 carbon atoms, preferably from 4 to 7 carbon atoms, and most preferably from 5 to 6 carbon atoms in the ring. Bicyclic carbocyclic aliphatic rings contain from 8 to 12 carbon atoms, preferably from 9 to 10 carbon atoms in the ring. Carbocyclic aliphatic rings may be unsubstituted or substituted with from 1 to 4 substituents on the ring. Preferred carbocyclic aliphatic ring substituents include: halo, cyano, alkyl, heteroalkyl, haloalkyl, phenyl, phenoxy or any combination thereof. More preferred substituents include halo and haloalkyl. Preferred carbocyclic aliphatic rings include cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, and cyclooctyl. More preferred carbocyclic aliphatic rings include cyclohexyl, cycloheptyl, and cyclooctyl. The most preferred carbocyclic aliphatic ring is cycloheptyl.

"Halo" is fluoro, chloro, bromo or iodo. Preferred halo are fluoro, chloro and bromo; more preferred are chloro and fluoro, especially fluoro.

"Haloalkyl" is a straight, branched, or cyclic hydrocarbon substituted with one or more halo substituents. Preferred haloalkyl are C₁-C₁₂; more preferred are C₁-C₆; more preferred still are C₁-C₃. Preferred halo substituents are fluoro and chloro. The most preferred haloalkyl is trifluoromethyl.

"Heteroalkyl" is a saturated or unsaturated chain containing carbon and at least one heteroatom, wherein no two heteroatoms are adjacent. Heteroalkyl chains contain from 1 to 18 member atoms (carbon and heteroatoms) in the chain, preferably 1 to 12, more preferably 1 to 6, more preferably still 1 to 4. Heteroalkyl chains may be straight or branched. Preferred branched heteroalkyl have one or two branches, preferably one branch. Preferred heteroalkyl are saturated. Unsaturated heteroalkyl have one or more double bonds and/or one or more triple bonds. Preferred unsaturated heteroalkyl have

one or two double bonds or one triple bond, more preferably one double bond. Heteroalkyl chains may be unsubstituted or substituted with from 1 to 4 substituents. Preferred heteroalkyl are unsubstituted. Preferred heteroalkyl substituents include halo, hydroxy, aryl (e.g., phenyl, tolyl, alkylphenyl, alkylphenyl, halophenyl), heterocyclyl, and heteroaryl. For example, alkyl substituted with the following substituents are heteroalkyl: alkoxy (e.g., methoxy, ethoxy, propoxy, butoxy, pentoxy), aryloxy (e.g., phenoxy, chlorophenoxy, tolyloxy, methoxyphenoxy, benzyloxy, alkylphenylphenoxy, acyloxyphenoxy), acyloxy (e.g., propionyloxy, benzoyloxy, acetoxyl), carbamoyloxy, carboxy, mercapto, alkylthio, acylthio, arylthio (e.g., phenylthio, chlorophenylthio, alkylphenylthio, alkoxyphenylthio, benzyloxyphenylthio, alkylphenylthio, alkoxyphenylthio, benzyloxyphenylthio), amino (e.g., amino, mono- and di- C₁-C₃ alkylamino, methylphenylamino, methylbenzylamino, C₁-C₃ alkylamido, carbamamido, ureido, guanidino).

"Heteroatom" is a nitrogen, sulfur, or oxygen atom. Groups containing more than one heteroatom may contain different heteroatoms.

"Heterocyclic aliphatic ring" is a saturated or unsaturated ring containing carbon and from 1 to about 4 heteroatoms in the ring, wherein no two heteroatoms are adjacent in the ring and no carbon in the ring that has a heteroatom attached to it also has a hydroxyl, amino, or thiol group attached to it. Heterocyclic aliphatic rings are not aromatic. Heterocyclic aliphatic rings are monocyclic, or are fused or bridged bicyclic ring systems. Monocyclic heterocyclic aliphatic rings contain from about 4 to about 10 member atoms (carbon and heteroatoms), preferably from 4 to 7, and most preferably from 5 to 6 in the ring. Bicyclic heterocyclic aliphatic rings contain from 8 to 12 member atoms, preferably 9 or 10 in the ring. Heterocyclic aliphatic rings may be unsubstituted or substituted with from 1 to 4 substituents on the ring. Preferred heterocyclic aliphatic ring substituents include: halo, cyano, alkyl, heteroalkyl, haloalkyl, phenyl, phenoxy or any combination thereof. More preferred substituents include halo and haloalkyl. Preferred heterocyclic aliphatic rings include piperzyl, morpholinyl, tetrahydrofuranyl, tetrahydropyranyl and piperdyl.

"Heteroaromatic ring" is an aromatic ring system containing carbon and from 1 to about 4 heteroatoms in the ring. Heteroaromatic rings are monocyclic or fused bicyclic ring systems. Monocyclic heteroaromatic rings contain from about 5 to about 10 member atoms (carbon and heteroatoms), preferably from 5 to 7, and most preferably from 5 to 6 in the ring. Bicyclic heteroaromatic rings contain from 8 to 12 member atoms, preferably 9 or 10 in the ring. Heteroaromatic rings may be unsubstituted or substituted with from 1 to 4 substituents on the ring. Preferred heteroaromatic ring

substituents include: halo, cyano, alkyl, heteroalkyl, haloalkyl, phenyl, phenoxy or any combination thereof. More preferred substituents include halo, haloalkyl, and phenyl. Preferred heteroaromatic rings include thienyl, thiazolo, purinyl, pyrimidyl, pyridyl, and furanyl. More preferred heteroaromatic rings include thienyl, furanyl, and pyridyl. The most preferred heteroaromatic ring is thienyl.


"Hydroxyalkyl" means HO-alkyl.

"Lower alkyl" is an alkyl chain radical comprised of 1 to 6, preferably 1 to 4 carbon atoms.

"Phenyl" is a monocyclic aromatic ring which may or may not be substituted with from about 1 to about 4 substituents. The substituents may be substituted at the *ortho*, *meta* or *para* position on the phenyl ring, or any combination thereof. Preferred phenyl substituents include: halo, cyano, alkyl, heteroalkyl, haloalkyl, phenyl, phenoxy or any combination thereof. More preferred substituents on the phenyl ring include halo and haloalkyl. The most preferred substituent is halo. The preferred substitution pattern on the phenyl ring is *ortho* or *meta*. The most preferred substitution pattern on the phenyl ring is *ortho*.

Compounds

The subject invention involves compounds having the following structure:



In the above structure, R_1 is CO_2H , $\text{C}(\text{O})\text{NHOH}$, CO_2R_5 , CH_2OH , $\text{S}(\text{O})_2\text{R}_5$, $\text{C}(\text{O})\text{NHR}_5$, $\text{C}(\text{O})\text{NHS}(\text{O})_2\text{R}_5$, or tetrazole; wherein R_5 is alkyl, heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring. Preferred R_5 is CH_3 , C_2H_5 , C_3H_7 . Preferred R_1 is CO_2H , $\text{C}(\text{O})\text{NHOH}$, CO_2CH_3 , $\text{CO}_2\text{C}_2\text{H}_5$, $\text{CO}_2\text{C}_3\text{H}_7$, $\text{CO}_2\text{C}_4\text{H}_9$, $\text{CO}_2\text{C}_3\text{H}_7\text{O}_2$, and $\text{C}(\text{O})\text{NHS}(\text{O})_2\text{R}_5$. More preferred R_1 is CO_2H , $\text{C}(\text{O})\text{NHOH}$, CO_2CH_3 , and $\text{CO}_2\text{C}_3\text{H}_5$. Most preferred R_1 is CO_2H and CO_2CH_3 .

In the above structure, R_2 is H or lower alkyl. Preferred R_2 is H and CH_3 . Most preferred R_2 is H.

In the above structure, X is NR_6R_7 , OR_8 , SR_9 , $\text{S}(\text{O})\text{R}_9$, $\text{S}(\text{O})_2\text{R}_9$, or F; wherein R_6 , R_7 and R_8 are selected independently from the group consisting of H, acyl, alkyl,

heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring; and wherein R_9 is alkyl, heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring. Preferred R_6 and R_7 are H, CH_3 and acyl. Preferred R_8 is H, CH_3 , CH_2CH_2OH . Preferred R_9 is CH_3 and CH_2CH_2OH . Preferred X is NR_6R_7 and OR_8 . Most preferred X is OH.

In the above structure, R_3 and R_4 are independently H, alkyl, hydroxyalkyl, alkoxyalkyl, OR_{10} , or SR_{10} , except that both R_3 and R_4 are not H; wherein R_{10} is alkyl, heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring. R_{10} has from 1 to about 8 member atoms. Preferred R_3 is H and CH_3 . Preferred R_4 is H and CH_3 . Preferred R_{10} is CH_3 .

In the above structure, Z is carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring. Preferred Z is monocyclic carbocyclic aliphatic ring, monocyclic heterocyclic aliphatic ring, monocyclic aromatic ring and monocyclic heteroaromatic ring. More preferred Z is monocyclic aromatic ring and monocyclic heteroaromatic ring. The most preferred Z is thienyl or phenyl.

The invention also includes optical isomers, diastereomers and enantiomers of the above structure. Thus, at all stereocenters where stereochemistry is not defined (C_{11} , C_{12} , C_{15} , and C_{16}), both epimers are envisioned. Preferred stereochemistry at all such stereocenters of the compounds of the invention mimic that of naturally occurring $PGF_{2\alpha}$.

It has been discovered that the novel PGF analogs of the subject invention are useful for treating bone disorders, especially those that require a significant increase in bone mass, bone volume, or bone strength. Surprisingly, the compounds of the subject invention have been found to provide the following advantages over known bone disorder therapies: (1) An increase trabecular number through formation of new trabeculae; (2) An increase in bone mass and bone volume while maintaining a more normal bone turnover rate; and (3) An increase in bone formation at the endosteal surface without increasing cortical porosity.

In order to determine and assess pharmacological activity, testing of the subject compounds in animals is carried out using various assays known to those skilled in the art. For example, the bone activity of the subject compounds can be conveniently demonstrated using an assay designed to test the ability of the subject compounds to increase bone volume, mass, or density. An example of such assays is the ovariectomized rat assay.


In the ovariectomized rat assay, six-month old rats are ovariectomized, aged 2 months, and then dosed once a day subcutaneously with a test compound. Upon

completion of the study, bone mass and/or density can be measured by dual energy x-ray absorptometry (DXA) or peripheral quantitative computed tomography (pQCT), or micro computed tomography (mCT). Alternatively, static and dynamic histomorphometry can be used to measure the increase in bone volume or formation.

Pharmacological activity for glaucoma can be demonstrated using assays designed to test the ability of the subject compounds to decrease intraocular pressure. Examples of such assays are described in the following reference, incorporated herein: C. Liljebris, G. Selen, B. Resul, J. Sternschantz, and U. Hacksell, "Derivatives of 17-Phenyl-18,19,20-trinorprostaglandin $F_{2\alpha}$ Isopropyl Ester: Potential Antiglaucoma Agents", Journal of Medicinal Chemistry, Vol. 38 No. 2 (1995), pp. 289-304.

Compounds useful in the subject invention can be made using conventional organic syntheses. Particularly preferred syntheses include the following two general reaction schemes:

Scheme 1



In **Scheme 1**, R_1 , R_2 , R_3 , R_4 , X , and Z are as defined above. The Methyl 7[3-(*R*)-hydroxy-5-oxo-1-cyclopent-1-yl] heptanoate (**S1a**) depicted as starting material for **Scheme 1** is commercially available (such as from the Sumitomo Chemical Company, Tokyo, Japan or the Cayman Chemical Company, Ann Arbor, MI).


In the above **Scheme 1**, Methyl 7[3-(*R*)-hydroxy-5-oxo-1-cyclopent-1-yl] heptanoate (**S1a**) is protected, preferably as a silyl ether, and is subjected to conditions which result in a 1,4 addition reaction of a suitably-substituted carbon nucleophile. For example, a cuprate addition of a homo-allyl alkene may be used. These manipulations result in ketones of the type depicted by **S1b**, which are readily reduced to alcohols, preferably with a boron reducing agent. These alcohols can be protected if needed during further manipulations. These manipulations are followed by the epoxidation of the olefin with a mild epoxidizing agent, such as meta chloroperbenzoic acid, to give compounds of the type depicted by **S1c**. These epoxides are reacted with oxygen nucleophiles under a great variety of conditions, but especially in the presence of Lewis

acids, to create compounds depicted by **Formula I**. Examples 4 and 5 illustrate compounds of **Formula I**.

Compounds depicted by **Formula II** are readily made from compounds of **Formula I** by their reaction with known ester manipulating reagents. For example, hydroxylamine transforms the ester functionality into a hydroxamic acid and LiOH transforms the ester to an acid. Examples 1-3 and 13-16 illustrate compounds of **Formula II**.

To create compounds depicted by **Formula III**, (Illustrated by Example 20) compounds are treated with a variety of agents known to transform secondary alcohols into halides, ethers, amines, sulfides and sulfoxides. Such reactions are disclosed, for example, in Advanced Organic Chemistry, by Jerry March. The compounds of **Formula III** may then be subsequently transformed at C₁ into the compounds depicted by **Formula II**, using substantially the same method for transforming the compounds of **Formula I** into the compounds depicted by **Formula II**.

Scheme 2



In **Scheme 2**, R_1 , R_2 , R_3 , R_4 , X , and Z are as defined above. The Corey lactone (**S2a**) depicted as starting material for **Scheme 2** is commercially available (such as from the Sumitomo Chemical Company, Tokyo, Japan; or the Cayman Chemical Company, Ann Arbor, MI).

Compounds depicted by **Formula IV** are made by elaboration of the commercially-available Corey lactone (**S2a**) to the advanced intermediate **S2b**. (An example of such a reaction is found in Example 6 herein). The intermediate **S2b** is coupled with beta-keto phosphonates, or the alcohols are suitably protected and olefination reactions are effected to give compounds of the type depicted by **S2c** after deprotection of the alcohol if appropriate. These compounds then undergo reduction of the alkenes and the ketone to give the compounds depicted by **Formula IV**. Example 12 illustrates a compound of **Formula IV**.

Compounds depicted by **Formula V** are derived from compounds of **S2c** by suitably protecting the alcohols, addition of R_2 , if desired, and reduction of the ketone and alkenes followed by suitable functional group manipulations as described above in **Scheme 1**. Examples 25, 26, and 28 illustrate compounds of **Formula V**.

Compounds depicted by **Formula VI** may be made from either compounds of **Formula IV** or **Formula V**, by reaction of the C_1 ester as described above in **Scheme 1**. Examples 6-11, 17 and 18, 22-24, 27, and 29-31 illustrate compounds of **Formula VI**.

These compounds are isolated by methods known to one of ordinary skill in the art. Such methods include, but are not limited to, extraction, solvent evaporation, distillation, and crystallization.

The following non-limiting examples illustrate the compounds, compositions, and uses of the present invention.

Examples


Compounds are analyzed using 1H and ^{13}C NMR, Elemental analysis, mass spectra, high resolution mass spectra and/or IR spectra as appropriate.

Typically, inert solvents are used, preferably in anhydrous form. For example, tetrahydrofuran (THF) is distilled from sodium and benzophenone, diisopropylamine is distilled from calcium hydride and all other solvents are purchased as the appropriate grade. Chromatography is performed on silica gel (70-230 mesh; Aldrich) or (230-400 mesh; Merck) as appropriate. Thin layer chromatography analysis is performed on glass mounted silica gel plates (200-300 mesh; Baker) and visualized using UV, 5%

phosphomolybdic acid in EtOH, or ammonium molybdate/ceric sulfate in 10% aqueous H₂SO₄.

Example 1

Preparation of 13,14-dihydro-16,16-dimethyl-16-(2-fluorophenoxy) 16-tetranor-prostaglandin F₁α:



a. Methyl 7-(2-oxo-4-(1,1,2,2-tetramethyl-1-silapropoxy)cyclopent-1-enyl) heptanoate: To a solution of Methyl-7-[3-(*R*)-hydroxy-5-oxo-1-cyclopenten-1-yl] heptanoate (1 equiv.) in CH₂Cl₂ at -78°C is added 2,6 lutidine (1.3 equiv.) dropwise over 15 minutes. The solution is kept at -78°C, and TBDMS Triflate (1.2 equiv.) in CH₂Cl₂ is added dropwise over 15 minutes. The reaction is warmed gradually to room temperature and stirred at room temperature for 15 hours. Aqueous 1% HCl is added until the solution is pH < 5.0 and the layers are separated. The water layer is back-extracted with CH₂Cl₂ and the organic layers are combined. The organic layer is washed with brine, dried (Na₂SO₄) and concentrated. The residue is distilled under vacuum (10 mm Hg) to provide the silyl ether.

b. Methyl 7-(5-(4-methyl-pent-3-enyl)-2-hydroxy-4-(1,1,2,2-tetramethyl-1-silapropoxy)cyclopentyl) heptanoate: To a slurry of Mg⁰ powder (2 equiv.) in THF at room temperature is added one crystal of I₂ and 1-bromoisoprene (2 equiv.) dropwise over 10 minutes. The reaction is exothermic as the addition continues. After the addition is complete, the reaction is refluxed for 3 hours and cooled to room temperature. The Grignard is diluted with THF and is added *via* cannula to a 3-necked flask equipped with mechanical stirring and is charged with CuBr as the dimethyl sulfide

adduct (DMS) (2 equiv.) in a 1:1 solution of THF/DMS at -78°C . After the addition of the Grignard reagent (~20 minutes), the reaction is stirred for 1 hour at -78°C . A solution of the ketone (1 equiv.) in THF is then added dropwise over 25 minutes. The reaction is stirred at -78°C for 15 minutes then allowed to warm slowly to room temperature over 2 hours. The reaction is quenched with aqueous NH_4Cl and the excess DMS allowed to evaporate. The reaction is partitioned between brine and CH_2Cl_2 and the layers are separated. The aqueous layer is back-extracted with CH_2Cl_2 and the organic layers are combined and dried (Na_2SO_4). The solvent is removed *in vacuo* and the residue chromatographed on SiO_2 (10 % hexane/EtOAc) to give the ketone as a clear oil.

The ketone is dissolved in MeOH and cooled to -40°C . Sodium borohydride (0.9 equiv.) is added portionwise over 10 minutes. After the addition is complete, the reaction is stirred for 13 hours at -40°C and then for 12 hours at -78°C . The reaction is quenched with water, partitioned between brine and CH_2Cl_2 and the layers separated. The aqueous layer is back-extracted with CH_2Cl_2 and the organic layers combined and dried (Na_2SO_4). The solvent is removed *in vacuo* and the residue is chromatographed on SiO_2 (30 % EtOAc/hexanes) to give the alcohol as a colorless oil.

c. Methyl 7-(2-hydroxy-5-(2-(3,3-dimethyl(2-oxiranyl)ethyl)-4-(1,1,2,2-tetramethyl-1-silapropoxy)cyclopentyl)heptanoate: The alcohol (1 equiv.) is dissolved in CH_2Cl_2 and cooled to 0°C . Sodium bicarbonate is added, followed by m-CPBA (57%-85% purity) (3 equiv.) portionwise over 15 minutes. After the addition is complete, the reaction is stirred for 20 hours at room temperature. The reaction is poured onto water, partitioned between brine and CH_2Cl_2 and the layers are separated. The aqueous layer is back-extracted with CH_2Cl_2 and the organic layers are combined and dried (Na_2SO_4). The solvent is removed *in vacuo* and the residue is chromatographed on SiO_2 (20% EtOAc/hexanes) to give the pair of epoxide diastereomers.

d. 13,14-dihydro-16,16 dimethyl, 16-(2-fluorophenoxy)-16- tetranor prostaglandin $\text{F}_{1\alpha}$ methyl ester: In a round bottom flask, epoxide (1 equiv.) and dry toluene are placed, the flask is cooled to 0°C and then treated with 2-fluorophenol (1.2 eq) and Alumina (1g per 100mg of phenol). The ice bath is removed and the reaction heated under nitrogen overnight. TLC is used to monitor the reaction. Excess fluorophenol is added if necessary. The reaction is quenched with brine, and is extracted with CH_2Cl_2 . The organic layer is washed three times with 1N HCl, brine, dried (Na_2SO_4), and is concentrated. Without further purification, to this crude reaction mixture is added CH_3CN and HF/Pyridine (0.1 mmol) while the flask is kept at 0°C .


After 3 hours. at 0°C, the reaction is quenched with saturated aqueous NaCl. The aqueous layer is extracted three times with CH₂Cl₂, the organic layers combined and washed three times with 1N HCl, brine, and dried (Na₂SO₄). After column chromatography (7:3, Hexane: Ethyl Acetate) the ester is obtained.




e. **13,14-dihydro-16,16 dimethyl, 16-(2-fluorophenoxy) tetranor prostaglandin F₁α:** To a round bottom flask, 13,14-dihydro-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor prostaglandin F₁α methyl ester and a 3:1 THF water mixture are added. The flask is cooled to 0°C, then an excess (2.5 equiv.) of lithium hydroxide is added, the ice bath removed, and the reaction is stirred at room temperature overnight. CH₂Cl₂ and saturated citric acid are added to the reaction mixture, the aqueous layer is washed 3 times with CH₂Cl₂, the organic layers are combined and is washed with brine, dried (Na₂SO₄), and chromatographed (CH₂Cl₂:methanol: acetic acid, 9.6: 0.4: 0.015). The acid 13,14-dihydro-16,16 dimethyl, 16-(2-fluorophenoxy) 16-tetranor prostaglandin F₁α is obtained.

Utilizing substantially the method of Example 1 (and using the appropriate starting materials), the following subject compounds of Examples 2-5 are obtained.

Example 2


13,14-dihydro-16,16,-dimethyl,16-(2-methylphenoxy)-16-tetranor Prostaglandin F₁α



Example 3**13,14-dihydro-16,16,-dimethyl, 16-(2,3 difluorophenoxy)-16- tetranor Prostaglandin****F₁α****Example 4****13,14-dihydro-16,16,-dimethyl, 16-(2,5 difluorophenoxy)16- tetranor Prostaglandin****F₁α methyl ester****Example 5****13,14-dihydro-16,16,-dimethyl, 16-(3-fluoro-5-trifluoromethyl phenoxy) 16-tetranor****Prostaglandin F₁α methyl ester**

Example 6

Preparation of 13, 14-dihydro- 16, 16-dimethyl, 16-(4-chlorophenoxy)- tetranor
Prostaglandin F₁α:



a. **6-(2,5-dioxolanyl)-7-hydroxy-2-oxabicyclo[3.3.0]octan-3-one:** In a round bottom flask equipped with a magnetic stirbar is placed 1,2-bis(trimethylsilyloxy)ethane (1.3 eq.) in CH₂Cl₂ containing trimethylsilyl-trifluoromethanesulfonate at -78°C. To this is added, within 20 minutes, a solution of 1 (1 eq) in CH₂Cl₂. The reaction is stirred for

1 hour at -78°C and then slowly warmed to 25°C for 1 hour. The reaction is quenched at 0°C with water, extracted with CH₂Cl₂ (3 times), dried (MgSO₄), and concentrated *in vacuo* to give crude 2. To a well stirred solution of crude 2 (1 eq) in methanol at 0°C is added a suspension of sodium methoxide (1.2 eq) in MeOH. The reaction is stirred at 0°C for 1 hour and then warmed to 25°C for 1 hour. The reaction is neutralized with acidic ion exchange resin which is washed thoroughly with MeOH (5x). The filtrate is concentrated *in vacuo* to give a syrup which is subjected to flash chromatography on silica gel eluting with 4:1 hexane : ethyl acetate and 2% MeOH in CH₂Cl₂ to give 3.

b. 6-(2,5 dioxolanyl)-2-oxa-7-(1,1,2,2-tetramethyl-1-silapropoxy) bicyclo [3.3.0] octan-3-one: In a round bottom flask with a magnetic stir bar, is stirred a solution of 3 (1 eq) in CH₂Cl₂. To this solution is added dropwise at -78°C 2,6-lutidine (1.9 eq) followed by TBDMSOTf (1.8 eq). The reaction is stirred for 30 minutes at -78°C and then warmed to 25°C overnight. The reaction is quenched with water. The organic layer is washed with water (3 times), dried (MgSO₄), and concentrated *in vacuo* to give a yellow oil which is subjected to flash chromatography on silica gel (hexanes then 1% MeOH in CH₂Cl₂). The product is then washed with 1N HCl (2 times), 0.1N HCl (2 times), water, and brine to give 4.

c. methyl 7-(5-(2,5-dioxolanyl)-2-hydroxy-4-(1,1,2,2-tetramethyl-(1-silapropoxy) cyclopentyl)hept-5-enoate: In a round bottom flask with a magnetic stir bar, is stirred a solution of 4 (1 eq) in dry toluene. To this solution, at -78°C, is slowly added DiBAL (1.24 eq). The reaction mixture is stirred for 2 hours and then warmed to 0°C. Saturated NH₄Cl is added to the reaction mixture which is then slowly warmed to 25°C. Diluted with water, the insoluble precipitate is removed by suction filtration and the solid is washed with EtOAc (2 times). The liquid phase is extracted with EtOAc (3 times) and the combined organic phase is dried (MgSO₄) and concentrated *in vacuo* to give the product, 5. The product, 5, must either be used immediately or stored at -70°C overnight.

To a suspension of (4-carboxybutyl)triphenylphosphonium bromide (2.2 eq) in THF at 0°C under N₂ is added dropwise a solution of KHMDS (0.5 M KHMDS in toluene, 4.4 eq). The resulting deep orange color reaction mixture is stirred for 1 hour at 25°C. To the reaction mixture above at -78°C is added a solution of 5 (1 eq) in THF. The reaction mixture is allowed to warm to 25°C overnight. The reaction is quenched with water at 0°C and the pH is adjusted to 3.5-4.0 with 1N HCl. The water phase is extracted with EtOAc (3 times) and the combined organic phase is dried (MgSO₄) and is concentrated *in vacuo* to give a syrup containing crude acid. To a well stirred solution of crude acid in ether and MeOH at 0°C is added TMS-diazomethane until the

reaction mixture keeps a light yellow color. The addition of 1 drop of glacial acetic acid and thin layer chromatography verifies that the reaction has gone to completion. The reaction solution is concentrated *in vacuo* and purified via flash chromatography on silica gel (30% EtOAc in hexanes) yielding **6**.

d. methyl 7-(2,4-dihydroxy-5-formylcyclopentyl)hept-5-enoate: In a round bottomed flask with a magnetic stir bar is placed an amount of the ketal, **6**. To this flask is added a sufficient amount of a mixture of 2 parts acetone 1 part 1N HCl to bring the ketal completely into solution. This material is stirred until, by TLC, the starting material is consumed, typically overnight. The crude mixture containing the product **7** is extracted with ether and the ether extract esterified *in situ* with, preferably, TMS-diazomethane. This allows for the formation of the product **8**, which is purified by column chromatography (30% EtOAc/hexanes) or is taken on without further purification.

e. 1-(dimethoxyphosphono)-3-(4-chlorophenoxy)-3-methyl butan-2-one: In a dry flask under N₂ is added THF (anhydrous) and Methyl dimethyl phosphonate (1 equiv.). The solution is cooled to -78°C and nBuLi solution (1.1 eq, of 2.5 M solution in hexanes) is added dropwise and then is stirred for 1 hour. The ester of 2-(4-chlorophenoxy)-2-methyl propionic acid is added dropwise in THF. This is stirred overnight and is allowed to warm to room temperature. The crude mixture is quenched with saturated NH₄Cl, then extracted with CH₂Cl₂ and purified by flash chromatography (5% MeOH in CH₂Cl₂) to yield **9**.

f. 16,16-dimethyl-16-(4-chlorophenoxy)-15-oxo-16-tetranor PGF_{2α} methyl ester: In a round bottom flask equipped with a magnetic stirbar is placed 3-(2,4-dichlorophenoxy)-dimethyl-2-oxo-propylphosphonate (1.65 eq) in DME and water (30:1). To this solution is added lithium bromide (2 eq), triethylamine (5.30 eq), and methyl 7-(2-formyl-3,5-dihydroxycyclopentyl)hept-5-enoate (1.0 eq). The solution is stirred at room temperature for 24 hours. Ether is added and the solution is washed once with 0.1 N HCl, and brine. The organic layer is dried (MgSO₄), filtered, and concentrated under reduced pressure. Purification is effected by silica gel column chromatography (methanol/ CH₂Cl₂ 1:50) to give 16,16-dimethyl-16-(4-chlorophenoxy)-15-oxo-16-tetranor PGF_{2α} methyl ester, **10**.


g. 13,14-dihydro-16,16-dimethyl-16-(4-chlorophenoxy)-16-tetranor PGF_{1α}: In a flame-dried round-bottomed flask equipped with a stir bar is placed **10** (1.0 equiv.), and cerium trichloride (1.05 eq) in methanol. The solution is stirred at room temperature for 5 minutes. The solution is cooled to -10°C and sodium borohydride (1.02 eq) in methanol is added. The solution is stirred at -10°C for 3 hours. The




mixture is treated with water and the pH brought to ~6 with 1 N HCl. The mixture is extracted twice with ethyl acetate, and the organic layers combined, dried (MgSO₄), filtered and concentrated under reduced pressure. Purification is effected by silica gel column chromatography (3% methanol in CH₂Cl₂ to 5% methanol in CH₂Cl₂) to give the 15-(*R*) alcohol and the 15-(*S*) alcohol. In a flame-dried round-bottomed flask equipped with a stir bar is placed one or the other of the epimeric alcohols, or a mixture thereof (1.0 eq) and palladium on carbon (10% Pd on C) in ethyl acetate. The heterogeneous mixture is treated with hydrogen gas for 18 hours. The mixture is then filtered through Celite and concentrated under reduced pressure to give the title saturated prostaglandin as the methyl ester. In a round-bottomed flask equipped with a stir bar is placed said ester (1.0 eq) and lithium hydroxide monohydrate (1.8 eq) in a 50/50 THF water solution. The mixture is stirred at room temperature for 6 hours and then diluted with water and acidified to pH ~2-3 with 1N HCl. The aqueous phase is extracted ~3 times with ethyl acetate and the organic layers combined. The combined organic layers are dried (MgSO₄), filtered, and concentrated under reduced pressure to yield 13,14-dihydro-16,16-dimethyl-16-(4-chlorophenoxy)-16-tetranor PGF₁α.





Utilizing substantially the method of Example 6 (and using the appropriate starting materials), the following subject compounds of Examples 7-14 are obtained.


Example 7

13,14-dihydro-16-methyl-16-(3-chlorophenoxy)-16-tetranor prostaglandin F₁α




Example 8**13,14-dihydro-16-ethyl-16-(2-methylphenoxy)-16-tetranor Prostaglandin F_{1α}****Example 9****13,14-dihydro-16-isopropyl 16-(2-fluorophenoxy)-16- tetranor Prostaglandin F_{1α}****Example 10****13,14-dihydro-16-(hydroxymethyl)-16-phenoxy-16-tetranor Prostaglandin F_{1α}**



Example 11**13,14-dihydro-16-methyl-16-(4-ethylphenoxy)-16-tetranor Prostaglandin F_{1α}****Example 12****13,14-dihydro-16-methyl-16-(3-chlorophenoxy)-16-tetranor prostaglandin F_{1α}
methyl ester****Example 13****13,14-dihydro-16-methyl-16-(4-phenylphenoxy)-16-tetranor prostaglandin F_{1α}
isopropyl ester****Example 14****13,14-dihydro-16,16-dimethyl-16-(4-phenoxyphenoxy)-16-tetranor prostaglandin
F_{1α} isopropyl ester**


Example 15**Preparation of 13,14-dihydro-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F_{1a} hydroxamic acid**

In a flame-dried round-bottomed flask equipped with a magnetic stir bar is placed 13,14-dihydro-16,16-dimethyl-16-(2-fluorophenoxy) -16- tetranor Prostaglandin F_{1a} -methyl ester (Example 1) (1.0 equiv.) in methanol. To this solution is added hydroxylamine in methanol (1.25 equiv.) and the solution is stirred for 18 hours. The solution is treated with 1N hydrochloric acid and extracted with ethyl acetate. The organic layer is washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The residue is purified by chromatography to give 13,14-dihydro-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F_{1a} hydroxamic acid.

The compounds of Examples 16-18 are prepared in a substantially similar manner as Example 15 using the appropriate hydroxylamine or sulfonamide and the intermediate corresponding to the appropriate example.


Example 16**13,14-dihydro-16-methyl-16-(3-chlorophenoxy) tetranor Prostaglandin F_{1a} 1-hydroxamic acid**



Example 17**13,14-dihydro-16-methoxymethyl-16-(2,3-difluorophenoxy)-16- tetranor****Prostaglandin F_{1a} 1-hydroxamic acid****Example 18****13,14-dihydro-16-methyl-16-(3-methoxyphenoxy) tetranor Prostaglandin F_{1a} 1-N-****methanesulfonamide**

Example 19**Preparation of 13,14-dihydro-15-fluoro-16-(2-fluorophenoxy) tetranor
Prostaglandin F_{1a} methyl ester**

The bis-silyl ether of Example 1 is treated with diethylaminosulfur trifluoride (as disclosed in the following references: Org. React. Vol. 35 (1988) p. 513; J. Org. Chem. Vol. 40 (1975) p. 574; and references cited therein.) to give 13,14-dihydro-15-fluoro-16-(2-fluorophenoxy) tetranor Prostaglandin F_{1a} methyl ester after deprotection as described in Example 1.

Utilizing substantially the method of Example 1 (and making the appropriate modification of R₂), the following subject compounds of Examples 20 and 21 are obtained.

Example 20**13,14-dihydro-15-methyl-16,16-dimethyl-16-(2-fluorophenoxy)-16- tetranor****Prostaglandin F_{1a}**


Example 21**13,14-dihydro-15-fluoro-16-(2,3-difluorophenoxy)-16- tetranor PGF_{1a} 1-hydroxamic acid****Example 22****Preparation of 13,14-dihydro-15-methylthio-15-dehydroxy-16-methyl-16-(2-methylphenoxy)-16- tetranor Prostaglandin F_{1a}**

The appropriate bis-silylated compound synthesized as described in Example 1 is treated sequentially with methanesulfonyl chloride (1.2 equiv.) and base (1.2 equiv.) (as disclosed in the following references: *J.C.S. Chem. Comm.* (1975) p. 658; *Tetrahedron Lett.* (1975) p. 3183; and references cited therein.) to generate the intermediate mesylate which is then treated immediately with nucleophiles (sodium thiomethoxide) (as disclosed in *Tetrahedron Lett.* Vol. 23 (1982) p. 3463 and references cited therein.) to give the protected thioalkyl ether. Subsequent deprotection as described in Example 1 provides 13,14-dihydro-15-methylthio-15-dehydroxy-16-methyl-16-(2-methylphenoxy)-16- tetranor Prostaglandin F_{1a}.

Utilizing substantially the method of Example 22, the following subject compounds of Examples 23-25 are obtained.


Example 23

13,14-dihydro-15-methylthio-15-dehydroxy-16-methyl-16-(2-methylphenoxy) 16-tetranor Prostaglandin F_{1a} 1-hydroxamic acid




Example 24


13,14-dihydro-15-methoxy-16,16 dimethyl-16-(2-fluorophenoxy) 16-tetranor Prostaglandin F_{1a} hydroxamic acid



Example 25



13,14-dihydro-15-(ethoxy)-15-dehydroxy-16,16-dimethyl-16-phenoxy 16-tetranor Prostaglandin F_{1a} methyl ester



Example 26**Preparation of 13,14-dihydro-15-sulfonylmethyl-15-dehydroxy-16-methyl-16-(2-methylphenoxy)-tetranor Prostaglandin F_{1a} methyl ester:**

The methyl ester of Example 22 is treated with the appropriate oxidizing agent as disclosed in the following references: Tetrahedron Lett. (1982) p. 3467; Prostaglandins Vol. 24 (1982) p. 801; Tetrahedron Lett. Vol. 23 (1982) p. 1023; and references cited therein.

Utilizing substantially the method of Example 26, the following subject compound of Example 27 is obtained.


Example 27**13,14-dihydro-15-sulfoxymethyl-15-dehydroxy-16-methyl-16-(2-methylphenoxy)-16-tetranor Prostaglandin F_{1a}****Example 28****Preparation of 13,14-dihydro- 15-methyl- 15-methylamino- 15-dehydroxy- 16, 16-dimethyl- 16- (2-fluorophenoxy)-16- tetranor Prostaglandin F_{1a} methyl ester**

The appropriate intermediate from Example 2 is condensed with N-methylamine to give the imine. Addition of the methyl cerium nucleophile (~1.5 equiv.) (for examples of cerium chloride-mediated nucleophilic addition see: J. Org. Chem. Vol. 49 (1984) p. 3904; J. Am. Chem. Soc. Vol. 111 (1989) p. 4392; and references cited therein) gives the aminomethyl derivative, which is then transformed into 13,14-dihydro- 15-methyl-15-methylamino- 15-dehydroxy- 16, 16-dimethyl- 16- (2-fluorophenoxy)-16- tetranor Prostaglandin F_{1a} methyl ester.

Utilizing substantially the method of Example 28, the following subject compounds of Examples 29-31 are obtained.


Example 29


13,14-dihydro-15-methyl-15-methylamino-15-dehydroxy-16-methyl-16-(2-methylphenoxy)- tetranor Prostaglandin F1_a 1-hydroxamic acid



Example 30

13,14-dihydro-15-methyl-15-(N,N-dimethylamino)-16-ethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F_{1a} isopropyl ester



Example 31**13,14-dihydro-16-ethyl-16-(2,6-difluorophenoxy)-16- tetranor Prostaglandin F_{1a}
glyceryl ester****Compositions**

Compositions of the subject invention comprise a safe and effective amount of the subject compounds, and a pharmaceutically-acceptable carrier. As used herein, "safe and effective amount" means an amount of a compound sufficient to significantly induce a positive modification in the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgment. A safe and effective amount of a compound will vary with the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the particular pharmaceutically-acceptable carrier utilized, and like factors within the knowledge and expertise of the attending physician.

In addition to the compound, the compositions of the subject invention contain a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier", as used herein, means one or more compatible solid or liquid filler diluents or encapsulating substances which are suitable for administration to a subject. The term "compatible", as used herein, means that the components of the composition are capable of being commingled with the compound, and with each other, in a manner such that there is no interaction which would substantially reduce the pharmaceutical efficacy of the composition under ordinary use situations. Pharmaceutically-acceptable carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the subject being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carriers or components thereof are sugars, such as lactose, glucose and sucrose; starches, such as cornstarch and potato starch; cellulose and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, cellulose acetate; powdered tragacanth; malt; gelatin; talc; solid lubricants, such as stearic acid, magnesium

stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerin, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers, such as the Tweens®; wetting agents such as sodium lauryl sulfate; coloring agents; flavoring agents, excipients; tableting agents; stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline; and phosphate buffer solutions.

The choice of a pharmaceutically-acceptable carrier to be used in conjunction with a compound is basically determined by the way the compound is to be administered. The compounds of the present invention may be administered systemically. Routes of administration include transdermal; oral; parenterally, including subcutaneous or intravenous injection; topical; and/or intranasal.

The appropriate amount of the compound to be used may be determined by routine experimentation with animal models. Such models include, but are not limited to the intact and ovariectomized rat models, the ferret, canine, and non human primate models as well as disuse models.

Preferred unit dosage forms for injection include sterile solutions of water, physiological saline, or mixtures thereof. The pH of said solutions should be adjusted to about 7.4. Suitable carriers for injection or surgical implants include hydrogels, controlled- or sustained release devices, polylactic acid, and collagen matrices.

Suitable pharmaceutically-acceptable carriers for topical application include those suited for use in lotions, creams, gels and the like. If the compound is to be administered perorally, the preferred unit dosage form is tablets, capsules and the like. The pharmaceutically-acceptable carriers suitable for the preparation of unit dosage forms for oral administration are well-known in the art. Their selection will depend on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of the subject invention, and can be made without difficulty by those skilled in the art.

Methods of Use

The compounds of the present invention are useful in treating many medical disorders, including for example, ocular disorders, hypertension, fertility control, nasal congestion, neurogenic bladder disorder, gastrointestinal disorders, dermatological disorders, and osteoporosis.

The compounds of the present invention are useful in increasing bone volume and trabecular number through formation of new trabeculae, increasing bone mass while maintaining a normalized bone turnover rate, and formation of bone at the

endosteal surface without removing bone from the existing cortex. Thus, these compounds are useful in the treatment and prevention of bone disorders.

The preferred routes of administration for treating bone disorders are transdermal and intranasal. Other preferred routes of administration include rectal, sublingual, and oral.

The dosage range of the compound for systemic administration is from about 0.01 to about 1000 $\mu\text{g/kg}$ body weight, preferably from about 0.1 to about 100 $\mu\text{g/kg}$ per body weight, most preferably from about 1 to about 50 $\mu\text{g/kg}$ body weight per day. The transdermal dosages will be designed to attain similar serum or plasma levels, based upon techniques known to those skilled in the art of pharmacokinetics and transdermal formulations. Plasma levels for systemic administration are expected to be in the range of 0.01 to 100 nanograms/ml, more preferably from 0.05 to 50 ng/ml, and most preferably from 0.1 to 10 ng/ml. While these dosages are based upon a daily administration rate, weekly or monthly accumulated dosages may also be used to calculate the clinical requirements.

Dosages may be varied based on the patient being treated, the condition being treated, the severity of the condition being treated, the route of administration, etc. to achieve the desired effect.

The compounds of the present invention are also useful in decreasing intraocular pressure. Thus, these compounds are useful in the treatment of glaucoma. The preferred route of administration for treating glaucoma is topically.

Composition and Method Examples

The following non-limiting examples illustrate the subject invention. The following composition and method examples do not limit the invention, but provide guidance to the skilled artisan to prepare and use the compounds, compositions and methods of the invention. In each case other compounds within the invention may be substituted for the example compound shown below with similar results. The skilled practitioner will appreciate that the examples provide guidance and may be varied based on the condition being treated and the patient.

Example A

Pharmaceutical compositions in the form of tablets are prepared by conventional methods, such as mixing and direct compaction, formulated as follows:

<u>Ingredient</u>	<u>Quantity (mg per tablet)</u>
Compound of Example 1	5
Microcrystalline Cellulose	100
Sodium Starch Glycollate	30
Magnesium Stearate	3

When administered orally once daily, the above composition substantially increases bone volume in a patient suffering from osteoporosis.

Example B

Pharmaceutical compositions in liquid form are prepared by conventional methods, formulated as follows:

<u>Ingredient</u>	<u>Quantity</u>
Compound of Example 1	5 mg
Phosphate buffered physiological saline	10 mL
Methyl Paraben	0.05 mL

When 1.0 mL of the above composition is administered subcutaneously once daily, the above composition substantially increases bone volume in a patient suffering from osteoporosis.

Example C

Topical pharmaceutical compositions for lowering intraocular pressure are prepared by conventional methods and formulated as follows:

<u>Ingredient</u>	<u>Amount (wt %)</u>
Compound of Example 14	0.004
Dextran 70	0.1
Hydroxypropyl methylcellulose	0.3
Sodium Chloride	0.77
Potassium chloride	0.12
Disodium EDTA (Edetate disodium)	0.05
Benzalkonium chloride	0.01
HCl and/or NaOH	pH 7.2-7.5


Purified water

q.s. to 100%

While particular embodiments of the subject invention have been described, it would be obvious to those skilled in the art that various changes and modifications to the compositions disclosed herein can be made without departing from the spirit and scope of the invention. It is intended to cover, in the appended claims, all such modifications that are within the scope of this invention.

What is claimed is:

1. A compound having the structure:



characterized in that

- (a) R_1 is CO_2H , C(O)NHOH , CO_2R_5 , CH_2OH , $\text{S(O)}_2\text{R}_5$, C(O)NHR_5 , $\text{C(O)NHS(O)}_2\text{R}_5$, or tetrazole; characterized in that R_5 is alkyl, heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring;
- (b) R_2 is H or lower alkyl;
- (c) X is NR_6R_7 , OR_8 , SR_9 , S(O)R_9 , $\text{S(O)}_2\text{R}_9$, or F; characterized in that R_6 , R_7 , and R_8 are independently selected from the group consisting of H, acyl, alkyl, heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring; and characterized in that R_9 is alkyl, heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring;
- (d) R_3 and R_4 are independently H, alkyl, hydroxyalkyl, alkoxyalkyl, OR_{10} , or SR_{10} , except that both R_3 and R_4 are not H; characterized in that R_{10} is alkyl, heteroalkyl, carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring, R_{10} having from 1 to about 8 member atoms;
- (e) Z is carbocyclic aliphatic ring, heterocyclic aliphatic ring, aromatic ring, or heteroaromatic ring; and

any optical isomer, diastereomer, enantiomer of the above structure, or a pharmaceutically-acceptable salt, or biohydrolyzable amide, ester, or imide thereof.

2. The compound according to Claim 1 characterized in that R_1 is selected from the group consisting of CO_2H , $C(O)NHOH$, CO_2CH_3 , and $CO_2C_3H_7$.
3. The compound according to Claim 2 characterized in that R_2 is H or CH_3 .
4. The compound according to Claim 3 characterized in that X is OH.
5. The compound according to Claim 4 characterized in that Z is thienyl or phenyl.
6. The compound according to any preceding claim characterized in that Z is substituted, said substituents being independently selected from the group consisting of halo, alkyl, haloalkyl, cyano, alkoxy, phenyl, and phenoxy.
7. The compound according to any preceding claim characterized in that Z is substituted, said substituents being independently selected from the group consisting of halo, alkyl, and phenyl.
8. The use of a compound according to any preceding claim in the manufacture of a medicament for treating a bone disorder in a human or other mammal.
9. The use of Claim 8 characterized in that said bone disorder is osteoporosis.

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07C405/00 //A61K31/557

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	US 5 703 108 A (ROSATI ROBERT L ET AL) 30 December 1997 see claims	1,8,9
X	US 4 621 100 A (LUND JOHN E ET AL) 4 November 1986	8,9
Y	see claims	1-9
X	US 4 024 179 A (BINDRA JASJIT S ET AL) 17 May 1977 cited in the application	1-7
Y	see claims	1-9
X	DE 24 60 990 A (HOECHST AG) 1 July 1976 cited in the application see claims	1-7
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 November 1998

Date of mailing of the international search report

03/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Berte, M

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 23225 A (ALCON LAB INC) 3 July 1997 see page 16, line 6 - page 17, line 3 -----	1-9

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
US 5703108	A	30-12-1997	NONE		
US 4621100	A	04-11-1986	JP 2016288 B		16-04-1990
			JP 58029710 A		22-02-1983
US 4024179	A	17-05-1977	US 4087604 A		02-05-1978
			AR 200585 A		22-11-1974
			AR 202931 A		31-07-1975
			AR 199966 A		08-10-1974
			AT 360672 B		26-01-1981
			AT 544576 A		15-06-1980
			AT 353285 B		12-11-1979
			AT 544676 A		15-04-1979
			AT 367034 B		25-05-1982
			AT 936973 A		15-10-1981
			AU 470375 B		11-03-1976
			AU 6224773 A		08-05-1975
			BE 806995 A		07-05-1974
			CA 1085831 A		16-09-1980
			CA 1088930 A		04-11-1980
			CA 1088931 A		04-11-1980
			CH 597176 A		31-03-1978
			CS 182793 B		31-05-1978
			CS 182794 B		31-05-1978
			CS 182791 B		31-05-1978
			DD 109212 A		20-10-1974
			DD 117233 A		05-01-1976
			DD 118856 A		20-03-1976
			DD 119411 A		20-04-1976
			DE 2355540 A		22-05-1974
			DE 2365322 A		07-11-1974
			DK 449778 A		10-10-1978
			DK 144247 B		25-01-1982
			FI 60389 B		30-09-1981
			FR 2205335 A		31-05-1974
			FR 2279729 A		20-02-1976
			FR 2283146 A		26-03-1976
			GB 1456514 A		24-11-1976
			GB 1456513 A		24-11-1976
			GB 1456512 A		24-11-1976

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4024179 A		HK 27979 A	11-05-1979
		HK 28079 A	11-05-1979
		HK 28179 A	11-05-1979
		IE 39538 B	08-11-1978
		IE 39539 B	08-11-1978
		IE 39537 B	08-11-1978
		IN 139384 A	12-06-1976
		JP 52053841 A	30-04-1977
		JP 52057147 A	11-05-1977
		JP 982264 C	27-12-1979
		JP 49093342 A	05-09-1974
		JP 54016491 B	22-06-1979
		KE 2930 A	30-03-1979
		KE 2931 A	30-03-1979
		KE 2932 A	30-03-1979
DE 2460990 A	01-07-1976	AU 8772775 A	23-06-1977
		BE 836950 A	22-06-1976
		DK 581075 A	22-06-1976
		FR 2294696 A	16-07-1976
		JP 51086449 A	29-07-1976
		LU 74076 A	11-11-1976
		NL 7514654 A	23-06-1976
		SE 7514447 A	22-06-1976
		ZA 7507872 A	24-11-1976
WO 9723225 A	03-07-1997	AU 7680096 A	17-07-1997